

SITE SYNOPSIS

SITE NAME: LOUGH DERG (SHANNON) SPA

SITE CODE: 004058

Lough Derg is the largest of the Shannon Lakes, being some 40 km long. Its maximum breadth across the Scarriff Bay -Youghal Bay transect is 13 km but for most of its length it is less than 5 km wide. The lake is relatively shallow at the northern end being mostly 6 m in depth but in the middle region it has an axial trench and descends to over 25 m in places. The narrow southern end of the lake has the greatest average depth, with a maximum of 34 m. The greater part of the lake lies on Carboniferous limestone but the narrow southern section is underlain by Silurian strata. Most of the lower part of the lake is enclosed by hills on both sides, the Slieve Aughty Mountains to the west and the Arra Mountains to the east. The northern end is bordered by relatively flat, agricultural country. The lake shows the high hardness levels and alkaline pH to be expected from its mainly limestone catchment basin, and it has most recently been classified as a mesotrophic system. The lake has many small islands, especially on its western and northern sides. The shoreline is often fringed with swamp vegetation. Aquatic vegetation includes a range of charophyte species, including the Red Data Book species, *Chara tomentosa*. The shoreline is often fringed by swamp vegetation, comprised of such species as Common Reed (*Phragmites australis*), Great Fen-sedge (*Cladium mariscus*) and Bottle Sedge (*Carex rostrata*).

Lough Derg is of importance for both breeding and wintering birds. The site supports a nationally important breeding colony of Common Tern (55 pairs recorded in 1995). Management of one of the islands used for nesting has increased the area of suitable habitat available and prevented nests being destroyed by fluctuating water levels. Large numbers of Black-headed Gull have traditionally bred on the many islands (2,176 pairs in 1985) but the recent status of this species is not known. A large Cormorant colony occurs in trees on the islands near Portumna - 167 nests were counted in 1995 and 122 in 1999. Lough Derg is also a noted breeding site for Great Crested Grebe (47 pairs in 1995) and Tufted Duck (326 individuals in late May 1995).

In winter, the lake is important for a range of waterfowl species, especially diving ducks, with nationally important populations of Tufted Duck (1,029), Goldeneye (215) and Mute Swan (235) - figures are average peaks for 4 of the 5 seasons 1995/96-1999/00. Other species which occur include Cormorant (120), Whooper Swan (18), Wigeon (272), Teal (342), Mallard (417), Pochard (61), Black-headed Gull (814), Coot (229), Lapwing (1,346) and Little Grebe (14). Lough Derg has traditionally been used by a relatively small flock of Greenland White-fronted Goose based in the Lough Derg-Lough Graney area and possibly further afield. The mean flock size for the 5 winters 1989/90-1993/94 was only 22, but few sightings have been made in recent years, In March 2004, however, c. 20 birds were observed in the Scarriff Bay area indicating that a flock may still be present in the area.

Lough Derg is of conservation interest for its fish and freshwater invertebrates. Lampreys, listed on Annex II of the E.U. Habitats Directive, are known to occur and the lake contains a landlocked population of Sea Lamprey (*Petromyzon marinus*). Brook Lamprey (*Lampetra planeri*) is known to be common in the lower River Shannon catchment where all three Irish Lamprey species breed. The endangered fish species Pollan (*Coregonus autumnalis pollan*) is recorded from Lough Derg, one of only four sites (L. Neagh, L. Erne, L. Ree and L. Derg) in which it occurs. Lough Derg is also a well-known fishing lake with a good Trout (*Salmo trutta*) fishery. Atlantic Salmon (*Salmo salar*) also use the lake as a spawning ground.

Lough Derg was classified as being strongly eutrophic in the early 1990s. Since 1997, a monitoring programme on the Shannon lakes has shown that the symptoms of eutrophication previously documented (i.e. high chlorophyll level and reduced water visibility) have been ameliorated significantly. These reductions have coincided with the invasion of the Shannon system by the Zebra Mussel (*Dreissena polymorpha*), a species which feeds on plankton, and also with measures to reduce phosphorus in sewage plants in the catchment. Enrichment of the lake, both by agricultural run-off and sewage, remains a threat and could affect the bird populations, especially the diving duck. Whilst the presence of Zebra Mussel in Lough Derg appears to have improved water quality in the lake, in the long-term this invasive bivalve may threaten the ecology of the lake. Recreational activities presently cause some disturbance to the birds and an increase in such activities would be of concern.

Lough Derg SPA is of high ornithological importance as it supports nationally important breeding populations of Common Tern, Cormorant, Great Crested Grebe, and probably Tufted Duck and Black-headed Gull. In winter, it has nationally important populations of Tufted Duck and Goldeneye, as well as a range of other species including Whooper Swan. The site is still used on occasions by Greenland White-fronted Goose. The presence of Common Tern, Whooper Swan and Greenland White-fronted Goose is of particular note as these are listed on Annex I of the E.U. Birds Directive.